



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Theorems on Cardinal Numbers.

BY A. N. WHITEHEAD.

In this paper it is proved ($\star 2$) that for cardinal numbers (finite or infinite) if $\alpha < \alpha'$ and $\beta < \beta'$, then $\alpha + \beta < \alpha' + \beta'$. It is already known (cf. $\star 4\cdot 37$ in my article on *Cardinal Numbers*, in this Journal, Vol. XXIV, 1902) that in this case $\alpha + \beta \leq \alpha' + \beta'$. Another form of $\star 2$ is $\star 1$, which is the most useful form for deductions. By the help of $\star 1$, it is proved ($\star 3$ and $\star 4$) that if n be a finite number and α_0 (as usual) the first infinite number, and if $n \times \beta = \alpha_0 \times \beta$, then $\beta = \alpha_0 \times \beta$. It is already known (cf. Zermelo, Gott. Nachricht., 1901) that if $\beta = n \times \beta$, then $\beta = \alpha_0 \times \beta$; and also it is obvious that if $\beta = \alpha_0 \times \beta$, then $\beta = n \times \beta$: then the present theorem completes this set of results.

Furthermore it is proved ($\star 6$) that if α is of the form $\alpha_0 \times \gamma$ [i. e. an $N\alpha_0$, cf. *Cardinal Numbers*, $\star 30$] and $\beta < \alpha$, then $\beta + \sigma = \alpha$ implies $\sigma = \alpha$. This proposition was assumed without proof in my article on *Cardinal Numbers* (cf. $\star 16\cdot 22$, for example). It is already known that $\beta + \alpha = \alpha$.

It will be noticed that a condition attaches itself to the hypothesis of $\star 1$, and thence to the hypotheses of the succeeding theorems, namely (stating it generally), that any two numbers less than the numbers considered, are such that one of them is either greater than, or equal to, or less than the other. I am not aware of any proof that one of these relations must hold for any two cardinal numbers; but classes of infinite cardinal numbers are known for which this condition is true.

The proofs are written in Peano's notation explained in the article on *Cardinal Numbers*, already cited; except that here \subset is used for "is contained in," as applied to classes, and \supset is used for "implies," as applied to propositions, instead of \supset for both these ideas. Also, in the proofs where a hypothesis is made to hold for the remainder of the proof "constr" for "constructive" is written after it.

- ★ 1 $\alpha, \beta, \alpha', \beta' \varepsilon Nc. \alpha + \beta = \alpha' + \beta' : \mu \leq \alpha. \nu \leq \beta. \supset_{\mu, \nu} \mu \leq \nu : \supset : \alpha \geq \alpha'. \vee. \beta \geq \beta',$
 $[R \varepsilon 1 \Rightarrow 1. u \cap v = \Lambda. u \cup v = \rho. u' \cap v' = \Lambda. u' \cup v' = \check{\rho}, \text{ constr (1)}$
 $S \varepsilon 1 \Rightarrow 1. \sigma \subset u \cap \rho v'. \check{\sigma} = v \cap \rho u', \text{ constr (2)}$
 $p = (u \cup v) - (\sigma \cup \check{\sigma}). \text{ constr (3)}$
 $R' = R_p \cup SR \cup \check{S}R, \text{ constr (4)}$
 $(4). \supset. R' \varepsilon 1 \Rightarrow 1. \check{\rho}' u = \check{\rho} (u - \sigma) \cup \check{\rho} (v \cap \rho u'), \text{ (5)}$
 $(5). \supset. u' \subset \check{\rho}' u. \supset. Nc' u' \leq Nc' u, \text{ (6)}$
 $(2). (6). \supset : Nc' (u \cap \rho v') \geq Nc' (v \cap \rho u'). \supset. Nc' u' \leq Nc' u. \text{ (7)}$
 $\text{Similarly } Nc' (v \cap \rho u') \geq Nc' (u \cap \rho v'). \supset. Nc' v' \leq Nc' v, \text{ (8)}$
 $(7). (8) : \mu \leq \alpha. \nu \leq \beta. \supset_{\mu, \nu} \mu \leq \nu : u \varepsilon \alpha. v \varepsilon \beta. u' \varepsilon \alpha'. v' \varepsilon \beta' : \supset : \text{Prop}] ,$
- ★ 2 $\alpha, \beta, \alpha', \beta' \varepsilon Nc. \alpha < \alpha'. \beta < \beta' : \mu \leq \alpha. \nu \leq \beta. \supset_{\mu, \nu} \mu \leq \nu : \supset.$
 $\alpha + \beta < \alpha' + \beta',$
 $[\star 1. \supset. \text{Prop.}]$
- ★ 3 $m, n \varepsilon Nc \text{ fin} - \iota 0. m \geq n. \beta \varepsilon Nc : \mu < \beta. \nu < \beta. \supset_{\mu, \nu} \mu \leq \nu :$
 $(m + n) \times \beta = \alpha_0 \times \beta : \supset. m \times \beta = \alpha_0 \times \beta.$
 $[\star 1. \text{Hyp.} \supset : m \times \beta + n \times \beta = \alpha_0 \times \beta + \alpha_0 \times \beta. \supset :$
 $m \times \beta \geq \alpha_0 \times \beta. \vee. n \times \beta \geq \alpha_0 \times \beta, \text{ (1)}$
 $l \varepsilon Nc \text{ fin}. l \times \beta \geq \alpha_0 \times \beta. \supset. l \times \beta = \alpha_0 \times \beta, \text{ (2)}$
 $m \geq n. \supset : n \times \beta = \alpha_0 \times \beta. \supset. m \times \beta = \alpha_0 \times \beta, \text{ (3)}$
 $(1). (2). (3). \supset. \text{Prop.}]$
- ★ 4 $n \varepsilon Nc \text{ fin} - \iota 0. \beta \varepsilon Nc : \mu < \beta. \nu < \beta. \supset_{\mu, \nu} \mu \leq \nu :$
 $n \times \beta = \alpha_0 \times \beta : \supset. \beta = \alpha_0 \times \beta.$
 $[\star 3. \supset. \text{Prop.}]$
- ★ 5 $m, n \varepsilon Nc \text{ fin} - \iota 0. m \circ' n. \beta \varepsilon Nc : \mu < \beta. \nu < \beta. \supset_{\mu, \nu} \mu \leq \nu :$
 $m \times \beta = n \times \beta : \supset. \beta = \alpha_0 \times \beta,$
 $[m \times \beta = n \times \beta. \supset. m \times \beta = \alpha_0 \times \beta. \text{ (1)}$
- Note: This theorem (1) is given by Zermelo, *Gott. Nachrichten*, 1901.
- (1). ★ 4. $\supset. \text{Prop.}]$
- ★ 6 $\alpha \varepsilon N\alpha_0 : \mu \leq \alpha. \nu \leq \alpha. \supset_{\mu, \nu} \mu \leq \nu : \beta < \alpha. \beta + \sigma = \alpha : \supset : \sigma = \alpha,$
 $[\text{Hyp. } \beta + \sigma = \alpha. \supset. \beta + \sigma = \alpha + \alpha, \text{ (1)}$
 $(1). \star 1. \beta < \alpha. \supset. \sigma \geq \alpha, \text{ (2)}$
 $\beta + \sigma = \alpha. \supset. \sigma \leq \alpha, \text{ (3)}$
 $(2). (3). \supset. \text{Prop.}]$